

# Journal of Physiology and Pathophysiology

Volume 7 Number 4 July, 2016

ISSN 2141-260X



*Academic  
Journals*

## ABOUT JPAP

**Journal of Physiology and Pathophysiology (JPAP)** is published monthly (one volume per year) by Academic Journals.

**Journal of Physiology and Pathophysiology (JPAP)** is an open access journal that provides rapid publication (monthly) of articles in all areas of the subject such as effect of different sitting posture on pulmonary function, study of anti-plasmodial and toxicological effects of methanolic bark extract of *Chrysophyllum albidum*, serum testosterone and morphology of the testes in wistar rats following chronic garlic feeding etc.

The Journal welcomes the submission of manuscripts that meet the general criteria of significance and scientific excellence. Papers will be published shortly after acceptance. All articles published in JPAP are peer-reviewed.

### Contact Us

Editorial Office: [jpap@academicjournals.org](mailto:jpap@academicjournals.org)

Help Desk: [helpdesk@academicjournals.org](mailto:helpdesk@academicjournals.org)

Website: <http://www.academicjournals.org/journal/JPAP>

Submit manuscript online <http://ms.academicjournals.me/>

## Editor

### **Dr. Kazi Mirajul Hoque**

*Scientist and Ramalingaswami Fellow,  
Molecular Pathophysiology Division,  
National Institute of Cholera and Enteric Diseases,  
P-33, CIT Road, Scheme-XM, Beliaghata,  
Kolkata-700010,  
INDIA...*

## Editorial Board Members

### **Dr. Yan-Qiu YU**

*Department of Pathophysiology  
Sri Ramachandra Deemed University,  
College of Basic Medical Sciences,  
China Medical University  
92.Beier street, Heping District,  
Shenyang, 110001,  
P. R. China.*

### **Dr. Wanzala Wycliffe**

*P.O. Box 11177-Tom Mboya-00400,  
Nairobi,  
Kenya...*

### **Venkateswara Rao Gogineni**

*Research Scientist  
University of Illinois College of Medicine at Peoria  
Department of Cancer Biology and Pharmacology  
One Illini Drive  
Peoria, IL 61605  
USA*

...

## Editorial Team

*Dr. Ajibola, Abdulwahid  
Olabisi Onabanjo University, Ago-Iwoye,  
Nigeria.*

### **Dr. Mostafa Azimzadeh**

*Dep. of Life Science Engineering,  
Faculty of New Sciences and Technologies,  
University of Tehran, Tehran,  
Iran..*

### **Dr. Birendra Nath Mallick**

*School of Life Sciences, Jawaharlal Nehru University  
New Delhi - 110 067,  
India.*

### **Dr. Ben-zhi CAI**

*Department of Pharmacology, Harbin medical university,  
Harbin, 150081,  
China.*

**Dr. Linda R. Chambliss.**

*4440 North Dromedary Rd  
Panjab University,  
Phoenix, Arizona 85018,  
USA.*

**Dr. Rajesh Chandra Sharma**

*301 Sai Mahal Co- Hsg. Society,  
Sector -3, plot no. 40 New Panvel, Navi  
Mumbai, Maharashtra-410205,  
India.*

**Dr. De-Yu Chen,**

*Shantou University Medical College (SUMC),  
22 Xinling Road, Shantou, Guangdong, P. R. China,  
515041,  
China.*

**Dr. Ayman El-Meghawry El-Kenawy Ali,**

*22 El-Galaa St., El-Mansoura, Egypt.*

**Dr. Karunakaran Ganesh Kumar,**

*Department of Metabolism and Aging, The Scripps  
Research Institute,  
130 Scripps Way, Jupiter, FL-33458,  
USA.*

**Dr. Farouk M. F. El-Sabban,**

*Department of Family Sciences, College for Women,  
Kuwait University, P.O. Box 5969, Safat 13060  
State of Kuwait.*

**Dr. Majid Hassanpour-Ezatti,**

*Dept. of Biology, Science School, Shahed Univ. Tehran,  
Iran.*

**Dr. Hedef Dhafir El-Yassin,**

*College of Medicine, University of Baghdad,  
Bab-Almuadam, Baghdad,  
Iraq.*

**Dr. Irawan Satriotomo,**

*Dept. Comparative Bioscience, University of Wisconsin  
2015 Linden Dr., Madison, WI 53706,  
USA.*

**Dr. Jingdong Li,**

*Institute of Cardiovascular Disease, Union Hospital,  
Tongji Medical College, Huazhong University of Science  
and Technology  
Wuhan, Hubei Province, 430022,  
P.R.of China.*

**Dr. Kazi Mirajul Hoque**

*3501 St. Paul Street, APT# 819, Baltimore, MD-21218,  
USA.*

**Dr. Tao Lianyuan**

*RM S209, QingNianGongYu, XinKai Road 80#,Dong Dan,  
Dong Cheng  
District, Beijing 100005,  
China.*

## ARTICLE

**Age-related effects of lead poisoning on sex hormones  
in adult male Wistar rats**

**23**

A. B. Daku, A. I. Salisu and A. B. El-ta'alu

*Full Length Research Paper*

## Age-related effects of lead poisoning on sex hormones in adult male Wistar rats

A. B. Daku<sup>1\*</sup>, A. I. Salisu<sup>2</sup> and A. B. El-ta'alu<sup>2</sup>

<sup>1</sup>Department of Human Physiology Faculty of Medicine, Federal University, Dutse, Nigeria.

<sup>2</sup>Department of Human Physiology Faculty of Basic Medical Sciences, Bayero University, Kano, Nigeria.

Received 9 December, 2015, Accepted 26 February, 2016

The World Health Organization (WHO) estimates that, about a quarter of the diseases facing mankind today occur due to prolonged exposure to environmental pollution, and that most of these environment-related diseases are however, not easily detected and may be acquired during childhood and manifested later in adulthood. The aim of this work was to determine the sub-chronic effects of lead poisoning on and some sex hormones, as well as age-related changes on Wistar Rats. Thirty (30) of 3-, 5-, and 7-months old age-groups male Wistar rats were selected; each age group was divided into experimental (lead fed) and control (distil water fed) groups. The levels of sex hormones (LH, FSH and Testestrone) were measured using ELISA method, while blood lead concentration was determined using the method of Atomic Absorption Spectrophotometer. There was a significant ( $P<0.05$ ) increase between experimental and control groups in blood lead concentration and insignificant ( $P>0.05$ ) increase in LH in both experimental and control groups. Significant ( $P<0.05$ ) decrease in body weight, testosterone levels, FSH was observed between experimental and control groups. The effect of lead (Pb) on testosterone levels was more pronounced in older animals. It was concluded that, ingestion of lead acetate has an age related effect on male sex hormones in older rats.

**Key words:** Sex hormones, lead, male, Wistar rats.

### INTRODUCTION

Over the last three decades, there has been increasing global concern over public health impacts attributed to direct and indirect environmental pollution, in particular, the global burden of disease. The World Health Organization (WHO) estimates that, about a quarter of the diseases facing mankind today occur due to prolonged exposure to environmental pollution (WHO, 2000). Most of these environment-related diseases are however, not easily detected and may be acquired during

childhood and manifested later in adulthood (CCNM (Calgary Centre for Naturopathic Medicine), 2014).

Lead metal may be found in dirt, dust, house hold utensils, dishes, furniture, leaded petrol, paints, ceramics, food cans, make-ups, traditional remedies, batteries, soil and water in varying degrees of concentration; and lead poisoning usually occurs from repeated exposure to small amounts (Agency For Toxic Substances and Disease Registry (ATSDR), 1999) The three major sites

\*Corresponding author. E-mail: [abdaku2@yahoo.com](mailto:abdaku2@yahoo.com). Tel: +2348036070418.

**Table 1.** Weight of rats before experiment.

Age of animals (months)	Weight (g)	
	Groups	
	Experimental	Control
3 month old	107.20±13.70	96.66±28.80
5 month old	136.20±17.99	110.98±18.62
7 month old	150.37±14.07	149.45±18.27

  

Weight Changes in rats after experiment			
Age of Animals (months)	Weight (g)		
	Groups		
	Experimental	Control	P-Value
3 month old	77.43±3.38 <sup>*a</sup>	130.66±28.80	0.000
5 month old	107.88±12.80 <sup>*a</sup>	150.60±6.65	0.001
7 month old	134.35±8.97 <sup>*a</sup>	165.62±17.80	0.040

of lead intoxication are kidney, blood, and the nervous system (Bersenyi', 2003), it is also a significant immunotoxin (Koller, 1990). Leaded petrol, as one of the metal's many sources in the environment, is a good indicator of reduction in exposure to lead (Landrigan et al., 2000).

There are many reports on lead toxicity and its deleterious effects in various species of animals as well studies on its pharmacokinetics and genotoxicity (Suradkar et al., 2009), but very few researchers try to correlate haemato-biochemical alterations of lead acetate in laboratory animals especially in rats.

Sources of food and feed contamination are soil, industrial polluted air and agricultural sources such as lead-containing pesticides, phosphate fertilizers and food processes (Bersenyi', 2003). Lead has become a regulatory concern and subject of much interest because of its widespread distribution in environment due to its continuous emission from industrial sources, automobile exhaust and its pharmacological behaviour to remain bound to mammalian tissues for a long time (Freeman, 1970). Human exposure has changed in origin but probably not in amount over the last half century; the quantity of lead used in the 20<sup>th</sup> century far exceeds the total consumed in all previous years (Fatima, et al. 2011). Such heavy uses have caused both local and global contamination of the air, dust, and soil (Elgohary, 2009).

There are reports on the effects of lead poisoning on various body systems, organs, and blood parameters but studies are limited on age- related effects of the heavy metal. The purpose of the study was to determine the age- related sub-chronic effects of lead poisoning on male sex hormones.

## MATERIALS AND METHODS

### Materials and chemicals

1. Accubind Elisa Microwells, Monobind Inc Lake Forest; CA 92630. USA Product Code: 625-300) were used to estimate LH

concentration in blood samples

2. Accubind Elisa Microwells, Monobind Inc Lake Forest, CA 92630. USA; Product Code: 425-300) were used to estimate FSH concentration in blood samples

3. Accubind Elisa Microwells, Monobind Inc Lake Forest, CA 92630. USA; Product Code: 3725-300) were used to estimate Testosterone concentration in the blood samples

4. Atomic Absorption Spectrophotometer (BUCK Scientific; model: 210 VGP, USA) were used to estimate blood lead concentration in blood samples.

### Procedure

Thirty (30) male Wistar rats were selected of three age-groups, of 3-, 5-, and 7-months old; each age-group was divided into experimental and control groups respectively. The experimental (n=15) animals were given lead acetate solution at 250 mg/kg body weight per day (Ambali et al., 2011) for 22 days, while the control (n=15) were placed on distilled water only. Both experimental and control animals were acclimatized for 7 days prior to commencement of experiment; animals were housed in metallic cages and given free access to chow and water. After the last day of lead treatment, rats were lulled to sleep by intravenous injection of 0.5cm<sup>3</sup> of 0.4% solution of sodium thiopental (Greene, 2002) and afterwards decapitated. Blood samples were collected and separated into two: first sample collected portion was centrifuged to get plasma which was used for the estimation of plasma FSH, LH, and testosterone concentrations using ELISA tests; the second blood sample was used for spectrophotometric analysis of blood lead concentration in blood samples, using Atomic Absorption Spectrophotometer (BUCK Scientific; model: 210 VGP, USA); weight of rats was measured before and after the experiment.

Data were collected and presented in tables of mean±SD and analyzed using student independent T-test to compare difference between experimental and control groups, while ANOVA was used to compare significant difference in experimental animals between the three age groups. All analyses were performed using Statistical Package for Social Science (SPSS) (Windows Evaluation Version 20, LEAD Technologies, USA) at p-value ≤ 0.05 (Table 1).

## RESULTS

Data with similar letter are not significantly (P > 0.05)

**Table 2.** Blood lead levels of rats after experiment.

Age of animals (months)	Lead levels ( $\mu\text{g/dl}$ )		
	Groups		
	Experimental	Control	P-Value
3 month old	46.00 $\pm$ 6.00 <sup>*a</sup>	14.56 $\pm$ 7.65	0.000
5 month old	46.75 $\pm$ 18.95 <sup>*a</sup>	18.00 $\pm$ 3.65	0.039
7 month old	50.75 $\pm$ 12.65 <sup>*a</sup>	17.60 $\pm$ 4.50	0.002

Data with similar letter are not significantly ( $P > 0.05$ ) different. \* $P < 0.05$  compared to control animals.

**Table 3.** Levels of luteinizing hormone after experiment in rats.

Age of animals (months)	LH Levels (mIU/ml)		
	Groups		
	Experimental	Control	P-Value
3 month old	5.000 $\pm$ 4.66 <sup>a</sup>	3.400 $\pm$ 2.03	0.405
5 month old	1.500 $\pm$ 0.58 <sup>b</sup>	1.000 $\pm$ 0.56	0.143
7 month old	1.000 $\pm$ 0.18 <sup>b</sup>	1.750 $\pm$ 0.96	0.135

Data with similar letter are not significantly ( $P > 0.05$ ) different. \* $P < 0.05$  compared to control animals.

**Table 4.** Levels of follicle stimulating hormone after experiment in rats.

Age of animals (months)	FSH Levels (mIU/ml)		
	Group		
	Experimental	Control	P-Value
3 month old	0.333 $\pm$ 0.58 <sup>*a</sup>	3.800 $\pm$ 2.76	0.035
5 month old	2.750 $\pm$ 0.50 <sup>a</sup>	0.250 $\pm$ 0.50	0.200
7 month old	0.0325 $\pm$ 0.39 <sup>a</sup>	0.100 $\pm$ 0.82	0.157

Data with similar letter are not significantly ( $P > 0.05$ ) different. \* $P < 0.05$  compared to control animals.

different \* $P < 0.05$  compared to control animals. Table 1 represents values of weight (g) changes in animals after experiment in different age groups. Significant difference ( $p < 0.05$ ) was recorded between experimental and control animals among all three age groups.

Significant changes in weights were recorded in all age groups: weight gain of 34.6g in normal control, while experimental had mean weight loss of 29.66g. Also significant ( $p \leq 0.05$ ) weights changes in 5-months old animals were also found with weight gain in normal controls (42.5 g). However, a loss in weight was recorded in experimental group (28.25 g).

Mean weight gain of 16.25 g was observed in normal control of 7-month old animals; though a loss was recorded in experimental animals (16.00 g). There was no significant ( $P > 0.05$ ) difference in weight loss between the three age groups.

Table 2 represents values of Blood lead concentration in different animal age groups. Significant differences ( $P$

$\leq 0.05$ ) were recorded in experimental compared to control groups.

While there was no significant ( $P > 0.05$ ) difference in Blood Lead levels between all age groups among experimental animals.

Table 3 represents values for levels of luteinizing hormone in different animal age groups. There was no significant ( $P > 0.05$ ) difference between experimental and control groups. A significant ( $P < 0.05$ ) difference in LH level between 3-months old compared to 5- and 7-months old experimental animals was recorded with more level among 3-months old animals

Table 4 shows levels of Follicle Stimulating Hormone in different animal age groups. There was significant ( $P < 0.05$ ) reduction in experimental group compared to control animals in 3-months old animals. No significant ( $P > 0.05$ ) difference in FSH levels between all age groups in experimental animals was observed.

Table 5 represents values of testosterone levels in

**Table 5.** Testosterone levels of rats after experiment.

Age of Animals (months)	Testosterone levels (ng/ml)		
	Groups		
	Experimental	Control	P-value
3-month old	0.110±0.03 <sup>a</sup>	0.656±0.87	0.265
5-month old	1.310±1.53 <sup>*ac</sup>	4.200±3.77	0.045
7-month old	0.523±0.26 <sup>*bc</sup>	4.235±5.45	0.040

Data with similar letter are not significantly ( $P > 0.05$ ) different. \* $P < 0.05$  compared to control animals.

different animal age groups. Significant ( $P < 0.05$ ) reduction in Testosterone Levels was recorded in experimental animals of 5- and 7-months old compared to control rats.

Also a significant ( $P < 0.05$ ) decrease in testosterone levels between 3-months old compared to 7- months old experimental animals was recorded with more decrease among 7-months old animals.

## DISCUSSION

This study supports other workers (Cezard and Haguener, 1992) who suggested that the weight loss may be attributed to loss of appetite and gastrointestinal disturbances; as well as to interruption in absorption and overall metabolism of feed nutrients (Marchlewicz et al., 2006). Even though there was no significant ( $P > 0.05$ ) difference between the three age groups, lead effect on weight loss was observed to be more pronounced in the younger 3-months old rats compared to the older ones.

Lead effect on reduction of body weight has been reported by other workers (Szymezak et al., 1983; Aseth et al., 1995; Teijon et al., 2006; Jegede et al., 2013). Such loss of body weight might also be due to interruption of absorption and metabolism of essential feed nutrients by lead acetate (Marija et al., 2004).

Absorption of lead presumably took place in all lead-treated groups across all ages, as all control groups had plasma lead level not more than 18 µg/dl. This shows the absence of contamination of animals in these control groups. After absorption into the blood, 95% of lead is bound to erythrocytes due to their high affinity for the metal. These cells contain the majority of lead found in blood stream, and the remaining part stays in plasma to be carried to other tissues; lead acetate and lead nitrate administration have been shown to increase blood lead level in experimental animals (Sokol, 1987; Ferguson et al., 1998; Sharma et al., 2011; Monira, et al. 2012;). Also there was no significant ( $P > 0.05$ ) difference in blood lead levels between age groups among experimental animals, suggesting same manner of lead absorption between the age groups.

There is no significant difference in LH levels between

the experimental and control rats. This is in accordance to Sokol (1987) and Fatima et al. (2011) that lead showed no significant effect on LH level, but contrary to findings of Camoratto (1990), Mukherjee and Mukhopadhyay (2009) and Taiwo et al. (2010), who reported that lead decreases LH concentration in experimental animals. While, an age-related difference among experimental animals was recorded with higher increase in LH among 3-months old animals.

Studies in rats have also demonstrated direct effect of lead on the testes and interference in the hypothalamic-pituitary axis (Saxena et al., 1989); disruption of the hypothalamic control of pituitary hormone secretion and in turn, spermatogenesis (Sokol, 1987); also gonadotoxic (Taiwo, et al. 2010).

Lead caused significant reduction in FSH level among 3-months old age group in this study. This is in conformity with Sokol (1987), but contrary to Raule and Morra (1952) and Petrusz (1979), who reported that lead increased FSH concentration. Increased concentration of FSH has also been reported following lead exposure in rats (Petrusz et al., 1979). Even though unchanged FSH concentrations in lead acetate treated mice were also reported (Pinon-Lataillade, et al. 1995), such differences in FSH secretion levels might relate to different lead levels and/or the duration of exposure among subjects (Mohsen et al., 2011).

Lead poisoning caused age-related reduction of testosterone level in this study with decrease more pronounced among 7-months old animals. This significant ( $P \leq 0.05$ ) reduction in testosterone levels is also in conformity with reports of Cullen et al. (1983), Sokol (1987), Camoratto (1990), Mukherjee and Mukhopadhyay (2009), Fatima et al. (2011) and Taiwo et al. (2010), who reported similar effects. The study by Fatima et al. (2011) supports the hypothesis that lead exerts its toxic effects on the hypothalamic or supra-hypothalamic sites, leaving an intact pituitary-testicular axis intact. In this study lead showed no significant effect on the LH levels, but significant decrease in testosterone. This suggests that lead exerts its effect more at the testicular level. This study also supports the finding of Sokol (1987), Wadi (1999) and Taiwo et al. (2010) who ascerted that lead targets the spermatogenesis and

sperms within the epididymis by producing reproductive toxicity rather than acting within the hypothalamic-pituitary-testicular (H-P-T) axis. They also suggested that the gonadotoxic effects of lead are on the intra-testicular sites with minimal effects on hormonal levels and no effect on extra testicular sites. Lead induced imbalances in the HPT hormonal axis, which causes pituitary cells to release inappropriate levels of LH and change the steroid negative feedback loop (Ronis, et al. 1996), usually at the hypothalamus level (Grattan, et al. 1996).

## Conclusion

Lead increases LH levels and significantly reduces FSH, testosterone levels in the experimental group and this effect on LH was more pronounced in younger age group, while the older age groups showed lower testosterone levels.

## Conflict of interest

The authors have not declared any conflict of interest

## REFERENCES

- Agency for Toxic Substances and Disease Registry [ATSDR] (1999). Toxicological profile for lead, Update. for ATSDR, U.S. Public Health Services, Atlanta, GA.
- Ambali SF, Angani M, Shittu M, Kawu MU (2011). Hematological changes induced by subchronic co-administration of chlorpyrifos and lead in Wistar rats: Alleviating effect of vitamin C. *Der Pharmacia Sinica* 2(2):276-284.
- Aseth J, Jacobsen D, Andersen O, Wickstron E (1995). Treatment of mercury and lead poisoning with dimercaptosuccinic acid (DMSA) and sodium dimercapto-propanesulfonate (DMPS). *Analyst* 120:853.
- Bersenyi A (2003). Study of toxic metals (Cd, Pb, Hg and Ni) in rabbits and broiler chickens. Budapest. pp. 1-89.
- Calgary Centre for Naturopathic Medicine (CCNM) (2014). Environmental Pollutants Detoxification. Available at: <http://www.calgarynaturopathic.com/Programs/EnviroDetox.aspx>
- Camoratto AM (1990). Inhibition of rat pituitary growth hormone release by subclinical levels of lead. *Toxicologist* 10:641.
- Cezard C, Haguenoer JM (1992). Toxicology of lead. Technique and documentation. Lavoisier. pp. 172-173.
- Cullen MR, Robins JM, Eskenazi B (1983). Adult organic lead intoxication: presentation of 31 new cases and review of the recent advances in literature. *Medicine* 62:47-221.
- Elgohary AA (2009). Prophylactic Effect of Angelica Archangelica Against Acute Lead Toxicity in Albino Rabbits. *Roman. J. Biophys.* 19(4):259-275.
- Fatima R, Umar AK, Ayub M, Shaukat S (2011). Protective Role of Ginger on Lead Induced Derangement In Plasma Testosterone and LH Levels of Male Sprague Dawley Rats. *J. Ayub. Med. Coll.* 23(4):24-27.
- Ferguson SA, Holson RR, Gazzara RA (1998). Minimal behavioral effects from moderate postnatal lead treatment in rats. *Neurotoxicol. Teratol.* 20:637-643.
- Grattan DR, Rocca MS, Sagrillo CA, McCarthy MM, Selmanoff M (1996). Antiandrogen microimplants into the rostral medial preoptic area decrease gamma-aminobutyric acidergic neuronal activity and increase luteinizing hormone secretion in the intact male rat. *Endocrinology* 137:4167-4173.
- Greene SA (2002). Veterinary anesthesia and pain management secrets. Henlay and Belfuc Inc, P.55; 266. In: Abbas Bubakar El-ta'alu, Alhassan Adamu Jibrin and Ibrahim Slayman. (2014). Effect of Mechanical Stretching of the Skin on Collagen Fibril Thermal Stability. *Niger. J. Basic Appl. Sci.* 2014 22(182):39-46.
- Jegade AI, Ajadi MB, Akinloye O (2013). Modulatory effects of kolaviron (Garcinia kola extract) on spermogram and reproductive system of adult male wistar rats in lead acetate induced toxicity. *J. Toxicol. Env. Health Sci.* 5(7):121-130.
- Koller LD (1990). The immunotoxic effects of lead in lead-exposed laboratory animals, Micronutrients and immune functions: Cytokines and metabolism. In Bendich A, Chandra RK (eds.). *Ann. New York Acad. Sci.* 587:160-167.
- Landrigan P, Boffetta P, Apostoli P (2000). Reproductive Toxicity And Carcinogenicity of Lead: A Critical Review. *Am. J. Ind. Med.* 38:231-243.
- Marchlewicz M, Wiszniewska B, Gonet B, Baranowska BI, Safranow K, Kolasa A, Glabowski W, Kurzawa R, Jakubowska K, Rac ME (2006). Increased Lipid Peroxidation and Ascorbic Acid Utilization in Testis and Epididymis of Mice Chronically Exposed to Lead. *Biometals.* pp. 13-14.
- Marija V, Piasek M, Blanusa M, Saric M, Juresa D, Kostial K (2004). Succimer treatment and calcium supplementation reduce tissue lead in suckling rats. *J. Appl. Toxicol.* 24:123-128.
- Mohsen V, Derek RS, Ping-Chi H (2011). Review article: How does lead induce male infertility. *Iran. J. Reprod. Med.* 9(1):1-8.
- Monira AA, Nermin ME, Hamdy T (2012). The Protective Role of Rosemary (*Rosmarinus officinalis*) in Lead Acetate Induced Toxicity in Rats. *J. Appl. Sci. Res.* 8(6):3071-3082.
- Mukherjee S, Mukhopadhyay PK (2009). Studies on lead and arsenic toxicity in male rat gonads and its protection by high dietary protein supplementation. *Al Ameen J. Med. Sci.* 2(1):73-77.
- Petrusz P, Weaver CM, Grant LD, Mushak P, Krigman MR (1979). Lead poisoning and reproduction: effects on pituitary and serum gonadotropins in neonatal rats. *Environ. Res.* 19:383-391.
- Pinon-Lataillade G, Thoreux-Manlay A, Coffigny H, Masse R, Soufir JC (1995). Reproductive toxicity of chronic lead exposure in male and female mice. *Human Exposure Toxicol.* 14:872-878.
- Raule A, Morra G (1952). Preliminary study on anterior pituitary gonadotropins in lead poisoning. *Med Lav.* 43:5-262.
- Ronis MJ, Badger TM, Shema SJ, Roberson PK, Shaikh F (1996). Reproductive toxicity and growth effects in rats exposed to lead at different periods during development. *Toxicol. Appl. Pharmacol.* 136:361-371.
- Saxena DK, Hussain T, Lal B, Chandra SV (1986). Lead induced testicular dysfunction in weaned rats. *Ind. Health* 24:105-109.
- Sharma S, Sharma V, Paliwal R, Pracheta (2011). Lead toxicity, oxidative damage and health implications. A review. *Int. J. Biotechnol. Mol. Biol. Res.* 2(13):215-221.
- Sokol RZ (1987). Hormonal Effects of Lead Acetate in the Male Rat: Mechanism of Action. *Biol. Reprod.* 37:1135-1138.
- Suradkar SG, Ghodasara DJ, Vihol P, Patel J, Jaiswal V, Prajapati KS (2009). Haemato-biochemical alterations induced by lead acetate toxicity in wistar rats. *Vet. World* 2(11):429-431.
- Szymezak J, Zechalko A, Biernat J (1983). Effect of fodder fat type on blood plasma lipids in rats intoxicated with lead. *Bromatol Chem. Toxicol.* 16:94-98.
- Taiwo AM, Ige SO, Babalola OO (2010). Assessments of Possible Gonadotoxic Effect of Lead on Experimental Male Rabbits. *Global Vet.* 5(5):282-286.
- Wadi SA, Ahmad G (1999). Effects of lead on male reproductive system in mice. *J. Toxicol. Environ. Health* 56(7):513-521.
- WHO (2000). Environmental pollution. [Internet Material]. Available at: [http://www.who.int/topics/environmental\\_pollution/en/](http://www.who.int/topics/environmental_pollution/en/)



# Journal of Physiology and Pathophysiology

## Related Journals Published by Academic Journals

- *Journal of Physiotherapy and Occupational Therapy*
- *Journal of Parasitology and Vector Biology*
- *Clinical Reviews and Opinions*
- *Journal of Clinical Pathology and Forensic Medicine*
- *Journal of Infectious Diseases and Immunity*
- *International Journal of Nutrition and Metabolism*
- *Journal of Dentistry and Oral Hygiene*

**academicJournals**